



**JOHN P. DRAWBERT, MD & TROY L. BERG, MD**  
Sports Medicine & Arthroscopy

4212 SOUTHTOWNE DRIVE  
EAU CLAIRE WI 54701  
TEL 715.832.1400

2501 COUNTY HWY I  
CHIPPEWA FALLS WI 54729  
TEL 715.723.8514

---

## **Post-op Instructions for Carpal Tunnel Release**

These instructions are to compliment the information given by the nursing staff and physical therapists. They cover many of the common questions.

### **Wound Care**

- Dressings are to be kept clean and dry. A small amount of clear drainage or bleeding is normal. If the ace wrap feels too tight, you may loosen it and re-wrap it.
- You may remove the ace bandage and dressing on the third day following surgery. Dressing may be removed all the way down to the stitches. Please leave the stitches intact. You may re-cover them with 4 x 4 gauze pads or a large Band-Aid.
- You may get the incision wet when showering 3 days after surgery. The shower should be brief and the wound patted dry with a clean towel. No baths or soaking the incision until 2-3 weeks after surgery and scabs are absent.
- If purulent drainage (thick white or greenish in color) is coming from the wound, or the wound has increasing redness, or if you are having a temperature of 101 or higher, please report these symptoms to your surgeon or the doctor on call.

### **Pain and Swelling**

- To lessen pain and swelling, ice your hand for the first 1-3 days after surgery. Put the ice in a zip lock bag or towel and place over the surgical site. We recommend 20 minutes, 4-5 times per day.
- Narcotic pain medication will be prescribed for use after you leave the hospital. Try to wean down as tolerated. These medications can cause constipation and you may want to use an over the counter stool softener. Tylenol products may be used instead of the prescribed pain medication. The prescribed narcotic medications should not be taken at the same time as plain Tylenol. You may supplement your pain medication by using ibuprofen or Aleve for any breakthrough pain. It can help to stagger your pain medication with ibuprofen or Aleve as needed. If a refill of medication is needed, please call the office during regular business hours, Monday-Friday 8:00 a.m. to 5:00 p.m. In general, refills will not be made after hours or on weekends, so please plan ahead.
- Swelling to some degree is common after surgery. To reduce swelling, elevation is very helpful. Elevate the hand above the heart level for the first 2-5 days after surgery. Elevation for 30 minutes every 2 hours is a good initial recommendation. Excessive pain and swelling should be reported to your surgeon.

**Exercises**

Gentle hand and wrist range of motion exercises can begin the day of surgery. After the bulky dressings have been removed on post-op day three, you may start hand-strengthening exercises by squeezing a soft ball.

**Driving**

To drive you must no longer be taking narcotic pain pills (plain Tylenol is allowed). Also, you must feel strong and alert.

**Follow-up**

Make sure an appointment has been scheduled for you at Chippewa Valley Orthopedics & Sports Medicine for approximately 1-2 weeks after surgery.

---

**John Drawbert, MD                      Troy Berg, M.D.**  
**Chippewa Valley Orthopedics & Sports Medicine**  
**(715) 832-1400**

Revised 10-07