

Dr. Troy Berg, Dr. Brent Carlson, Dr. Nathan Harris
 Chippewa Valley Orthopedics & Sports Medicine
 1200 OakLeaf Way, Suite A 757 Lakeland Drive, Suite B
 Altoona, WI 54720 Chippewa Falls, WI 54729

Patient _____
 D.O.S. _____

SUBACROMIAL DECOMPRESSION PROTOCOL

	WEEK 1-2 Begins on DOS	WEEK 3-6	WEEK 7-12	WEEK 13+
PASSIVE SCAPTION	Painfree to full	Full		
ACTIVE SCAPTION	Start week 2 as tolerated	Painfree		
PASSIVE ER/IR	Start at 30° abd, progress to 45° abd	Full		
ACTIVE ER	Start week 2 as tolerated	Painfree		
IMMOB/SLING	As needed	No		
P.T. visits/week	1-3	1-3	1-3	1-3
EXERCISES				
<ul style="list-style-type: none"> • Individual exercise progression may vary <p>Questions? Please contact: Northwoods Therapy Associates</p> <p>Altoona, WI (715) 839-9266</p> <p>Chippewa Falls, WI (715) 723-5060</p> <p>Reviewed September 2015</p>	Passive Pendulums	Initiate isotonic Scaption, rowing, Sidelying ER, Abd to 90, Ext to neutral	Continue with all previous isotonic, strengthening, neuromuscular control and endurance exercises	Continue ROM and strengthening program and self capsular stretches as needed. Continue fundamental shoulder exercise program. Gradually return to overhead activities and progress as tolerated.
	Pulleys			
	Wand ER, IR, Scaption	Add 1# after one week of proper form and no pain	2 hand drills: Chest pass Side to side throws	
	Isometrics Flex, Ext, Abd, IR, ER, Biceps		At 9 weeks add one hand plyo's	
	Rhythmic Stab		Wall dribbles, baseball throws	
	ER/IR Tubing	Joint mobs	Shovel throws	
	Prone rowing	Scapular stabilization	Initiate sport program	
Goals: non-painful arc of motion. Prevent muscular atrophy, re-establish dynamic stabilization and decrease pain and inflammation. Criteria to advance: Full ROM, Minimal pain and tenderness and good MMT of IR, ER and flexion.	UE endurance activity	Goals: Improve strength, power and endurance. Improve neuromuscular control and prepare athlete to return to sport and worker back to work. Criteria to advance: Full ROM, no pain or tenderness, satisfactory muscular strength and satisfactory clinical exam.	Goals: Progressively increase activities to prepare for full functional return.	