



www.cvosm.com

JOHN C. BERSCHBACK, MD
Hand & Orthopedic Surgery

1200 OAKLEAF WAY STE A
ALTOONA WI 54720
TEL 715.832.1400

757 LAKELAND DR. STE B
CHIPPEWA FALLS WI 54729
TEL 715.723.8514

Post-op Instructions for Trigger Finger Release

These instructions are to compliment the information given by the nursing staff and Dr. Berschback. They cover many of the common questions.

WOUND CARE

Dressings are to be kept clean and dry. A small amount of clear drainage or bleeding is normal. If the ace wrap feels too tight, you may loosen it and re-wrap it.

You may remove the ace bandage and dressing on the second day following surgery. The dressing may be removed all the way down to the stitches. Please leave the stitches intact. You may recover them with 4x4 gauze pads and tape or a large Band-Aid.

You may get the incision wet when showering 2 days after surgery. The shower should be brief and the wound patted dry with a clean towel. No baths or soaking the incision until your stitches have been removed.

If purulent drainage (thick white or greenish in color) is coming from the wound, or the wound has increasing redness, or if you are having a temperature of 101 or higher, please report these symptoms to your surgeon or the doctor on call.

PAIN & SWELLING

To lessen pain and swelling, you may ice your hand for the first 1-3 days after surgery. Put the ice in a zip lock bag or towel and place over the surgical site. We recommend 20 minutes, 2-3 times per day.

Narcotic pain medication may be prescribed for use after you leave the hospital. Try to wean down as tolerated. These medications can cause constipation and you may want to use an over the counter stool softener. Tylenol products or medications such as Aleve or Motrin (anti-inflammatory) may be used instead of the prescribed pain medication, and this is encouraged if you are not having much pain. The prescribed narcotic medications should not be taken at the same time as plain Tylenol. If a refill of medication is needed, please call the office during regular business hours, Monday-Friday 8:00 a.m. to 5:00 p.m. In general, refills will not be made after hours or on weekends, so please plan ahead.

Swelling to some degree is common after surgery. To reduce swelling, elevation is very helpful. Elevate the hand above the heart level for the first few days after surgery. Elevation for 30 minutes every 2-3 hours is a good initial recommendation. Excessive pain and swelling should be reported to your surgeon. When you sleep, try to keep the hand elevated on a pillow.

EXERCISES

Gentle finger and wrist range of motion exercises can begin the day of surgery. You may continue to advance your activities as tolerated, but you should avoid heavy lifting.

DRIVING

To drive you must no longer be taking narcotic pain pills (plain Tylenol is allowed). Also, you must feel strong and alert.

FOLLOW-UP

Make sure an appointment has been scheduled for you at Chippewa Valley Orthopedics & Sports Medicine for approximately 2 weeks after surgery.

John Berschback, MD

Chippewa Valley Orthopedics & Sports Medicine

(715) 832-1400