

CHIPPEWA FALLS
ORTHOPEDICS AND
SPORTS MEDICINE

NATHAN H. HARRIS, MD
Orthopedic Surgeon

1200 OAKLEAF WAY STE A ALTOONA WI 54720
TEL 715.832.1400 | 800.322.1747 | FAX 715.832.4187

757 LAKELAND DR. STE B CHIPPEWA FALLS WI 54729
TEL 715.723.8514 | 800.322.1748 | FAX 715.723.5989

2200 CEDARCREST DR STE A RICE LAKE WI 54868
TEL 715.537.4550 | 800.322.1747 | FAX 715.832.4187

Rehabilitation Protocol

Distal Humerus Fracture Fixation

Phase I: Early ROM & Protect Fixation (0 to 4 weeks)

- Splint and postop dressing remains in place for the first 2 weeks.
- Hinged Elbow Brace worn at all times (after the initial dressing removed)
- Initiate elbow exercise program 5 times per day:
 - Passive extension to 0 degrees as tolerated
 - Active flexion to 90 as tolerated
 - Progress to full Supination/Pronation
- Shoulder/wrist/hand exercises encouraged
- Edema control
- Avoid active elbow extension x 6 wks postop
- Grip and wrist/hand AROM immediately.
- Perform supination ROM only with the elbow flexed to 90 degrees.

Phase II: Advance ROM and function (4-6 weeks)


- Passive extension to 0 degrees as tolerated
- Active/Passive flexion may increase by 10 degrees per week as tolerated
- Progress to full Supination/Pronation
- Progress to active/active assist ROM
- Begin wrist, forearm, and hand/finger strengthening exercises

Phase III: Full ROM and function (6-10 weeks)

- Wean from brace once fracture healing on XRAY (6-8 wks)
- May begin static progressive splinting if failure to achieve >100 degree arc ROM by 8 wks
- Begin active elbow extension
- Progress with strengthening slowly
 - -5 lbs lifting restriction at 6wks
 - -15 lbs lifting restriction at 8 wks
- Transition to HEP by 8-10 wks

Phase IV: Full ROM and function (10-12+ weeks)

- Gradually return to light activity and exercise
- No further activity restrictions after 12 weeks



NATHAN H. HARRIS, MD
Orthopedic Surgeon

1200 OAKLEAF WAY STE A ALTOONA WI 54720
TEL 715.832.1400 | 800.322.1747 | FAX 715.832.4187
757 LAKELAND DR. STE B CHIPPEVA FALLS WI 54729
TEL 715.723.8514 | 800.322.1748 | FAX 715.723.5989
2200 CEDARCREST DR STE A RICE LAKE WI 54868
TEL 715.537.4550 | 800.322.1747 | FAX 715.832.4187

Rehabilitation Protocol Distal Humerus Fracture Fixation

Phase I: Early ROM & Protect Fixation (0 to 4 weeks)

- Splint and postop dressing remains in place for the first 2 weeks.
- Hinged Elbow Brace worn at all times (after the initial dressing removed)
- Initiate elbow exercise program 5 times per day:
 - Passive extension to 0 degrees as tolerated
 - Active flexion to 90 as tolerated
 - Progress to full Supination/Pronation
- Shoulder/wrist/hand exercises encouraged
- Edema control
- Avoid active elbow extension x 6 wks postop
- Grip and wrist/hand AROM immediately.
- Perform supination ROM only with the elbow flexed to 90 degrees.

Phase II: Advance ROM and function (4-6 weeks)

- Passive extension to 0 degrees as tolerated
- Active/Passive flexion may increase by 10 degrees per week as tolerated
- Progress to full Supination/Pronation
- Progress to active/active assist ROM
- Begin wrist, forearm, and hand/finger strengthening exercises

Phase III: Full ROM and function (6-10 weeks)

- Wean from brace once fracture healing on XRAY (6-8 wks)
- May begin static progressive splinting if failure to achieve >100 degree arc ROM by 8 wks
- Begin active elbow extension
- Progress with strengthening slowly
 - -5 lbs lifting restriction at 6wks
 - -15 lbs lifting restriction at 8 wks
- Transition to HEP by 8-10 wks

Phase IV: Full ROM and function (10-12+ weeks)

- Gradually return to light activity and exercise
- No further activity restrictions after 12 weeks