

Post-op Instructions for Hardware Removal

These instructions are to compliment the information given by the nursing staff and physical therapists. They cover many common questions.

WOUND CARE

Dressings are to be kept clean and dry. A small amount of clear drainage or bleeding is normal. If an ace wrap is present and feels too tight, you may loosen it and re-wrap it.

If a splint is present, it should remain on until you follow-up in the clinic with your surgeon. When showering, please place a plastic bag over the splint and secure it with tape. You may wrap your arm with Glad Press & Seal plastic wrap instead of a plastic bag. To keep the splint dry, you may wrap a small towel around the splint prior to applying the plastic bag or wrap. Pat the splint dry immediately with a clean towel if it does get slightly wet.

If no splint is present, you may get the incision wet when showering 3 days after surgery. The shower should be brief and the wound patted dry with a clean towel. No baths or soaking the incision until 2 weeks after surgery and scabs are absent.

If purulent drainage (thick white or greenish in color) is coming from the wound, or the wound has increasing redness, or if you are having a temperature of 101 or higher, please report these symptoms to your surgeon or the doctor on call.

PAIN

To lessen pain and swelling, you may ice after surgery. Put the ice in a zip lock bag or towel and place over the surgical site. We recommend 20 minutes, 4-5 times per day for the first 1-3 days. Do not place ice or cooling devices directly on the skin for prolonged periods of time as it may damage the skin.

Narcotic pain medication will be prescribed for use after you leave the hospital. Try to wean down as tolerated. These medications can cause constipation and you may want to use an over the counter stool softener. Tylenol products may be used instead of the prescribed pain medication. The prescribed narcotic medications should not be taken at the same time as plain Tylenol. You may supplement your pain medication by using Ibuprofen or Aleve for any breakthrough pain. It can help to stagger your pain medication with Ibuprofen or Aleve as needed. If a refill medication is needed, please call the office during regular business hours, Monday - Friday 8:00 a.m. to 5:00 p.m. In general, refills will not be made after hours or on weekends, so please plan ahead.

Swelling to some degree is common after surgery. To reduce swelling, elevation is very helpful. Elevate the affected extremity above the heart level for the first 2-5 days after surgery. Elevation for 30 minutes every 2 hours is a good initial recommendation. Excessive pain and swelling should be reported to your surgeon.

DRIVING

To drive you must no longer be taking narcotic pain pills (plain Tylenol is allowed). Also, you must feel strong and alert.

Nathan Harris, MD
Chippewa Valley Orthopedics
715-832-1400